

Quality ID #205 (NQF 0409): HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis – National Quality Strategy Domain: Effective Clinical Care

2018 OPTIONS FOR INDIVIDUAL MEASURES:
REGISTRY ONLY

MEASURE TYPE:
Process

DESCRIPTION:
Percentage of patients aged 13 years and older with a diagnosis of HIV/AIDS for whom chlamydia, gonorrhea, and syphilis screenings were performed at least once since the diagnosis of HIV infection

INSTRUCTIONS:
This measure is to be submitted a minimum of **once per performance period** for patients with HIV/AIDS seen during the performance period. Only patients who had at least two visits during the performance period, with at least 90 days between each visit will be counted in the denominator for this measure. This measure is intended to reflect the quality of services provided for the primary management of patients with HIV/AIDS.

Measure Submission:
The listed denominator criteria is used to identify the intended patient population. The numerator options included in this specification are used to submit the quality actions allowed by the measure. The quality-data codes listed do not need to be submitted for registry-based submissions; however, these codes may be submitted for those registries that utilize claims data.

DENOMINATOR:
Patients aged 13 and older with a diagnosis of HIV/AIDS who had at least two medical visits during the measurement year, with at least 90 days between each visit

Denominator Criteria (Eligible Cases):

Patients aged ≥ 13 years of age on date of encounter

AND

Diagnosis for HIV/AIDS (ICD-10-CM): Z21, B20

AND

Patient encounters during the performance period (CPT or HCPCS): 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, G0402

AND

At Least Two Denominator Eligible Encounters During the Measurement Year, With at Least 90 days Between Each

AND NOT

DENOMINATOR EXCLUSION:

Patients who use hospice services any time during the measurement period: G9725

NUMERATOR:
Patients with chlamydia, gonorrhea, and syphilis screenings performed at least once since the diagnosis of HIV infection

NUMERATOR NOTE: *Submit G9228 when results are documented for all of the 3 screenings*

Numerator Options:
Performance Met:

Chlamydia, gonorrhea and syphilis screening results documented (report when results are present for all of the 3 screenings) (G9228)

OR

Denominator Exception:

Chlamydia, gonorrhea, and syphilis screening results not documented (Patient refusal is the only allowed exception) (G9229)

OR

Performance Not Met:

Chlamydia, gonorrhea, and syphilis screening results not documented as performed, reason not given (G9230)

RATIONALE:

Sexually transmitted diseases that cause mucosal inflammation (such as gonorrhea and chlamydia) increase the risk for HIV-infection (as these diseases and other sexually transmitted diseases can increase the infectiousness of and a person's susceptibility to HIV) (Galvin, 2004).

CLINICAL RECOMMENDATION STATEMENTS:

All patients should be screened with laboratory tests for STDs at the initial encounter (A-II for syphilis, for trichomoniasis in women, and for chlamydial infection in women aged less than 25 years; B-II for gonorrhea and chlamydial infection in all men and women), and thereafter, depending on reported high-risk behavior, the presence of other STDs, and the prevalence of STDs in the community (B-III). (Aberg, 2004)

Consideration should be given to screening all HIV-infected men and women for gonorrhea and chlamydial infections. However, because of the cost of screening and the variability of prevalence of these infections, decisions about routine screening for these infections should be based on epidemiologic factors (including prevalence of infection in the community or the population being served), availability of tests, and cost. (Some HIV specialists also recommend type-specific serologic testing for herpes simplex virus type 2 for both men and women.) (B-II, for identifying STDs) (CDC, HRSA, NIH, HIVMA of IDSA, 2003)

COPYRIGHT:


This Physician Performance Measure (Measure) and related data specifications have been developed by the PCPI(R) Foundation (PCPI[R]) and the National Committee for Quality Assurance (NCQA). This Measure is not a clinical guideline and does not establish a standard of medical care, and has not been tested for all potential applications. The Measure, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, eg, use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the Measure for commercial gain, or incorporation of the Measure into a product or service that is sold, licensed or distributed for commercial gain. Commercial uses of the Measure require a license agreement between the user and the PCPI(R) or NCQA. Neither the American Medical Association (AMA), nor the former AMA-convened Physician Consortium for Performance Improvement(R), PCPI, NCQA nor its members shall be responsible for any use of the Measure. (C) 2017 National Committee for Quality Assurance and PCPI (R) Foundation. All Rights Reserved.

Limited proprietary coding is contained in the Measure specifications for user convenience. Users of proprietary code sets should obtain all necessary licenses from the owners of the code sets. NCQA disclaims all liability for use or accuracy of any CPT or other codes contained in the specifications.

CPT(R) contained in the Measure specifications is copyright 2004-2017 American Medical Association. LOINC(R) copyright 2004-2017 Regenstrief Institute, Inc. This material contains SNOMED Clinical Terms(R) (SNOMED CT[R]) copyright 2004-2017 International Health Terminology Standards Development Organisation. ICD-10 copyright 2017 World Health Organization. All Rights Reserved.

The performance Measure is not a clinical guideline and does not establish a standard of medical care, and has not been tested for all potential applications. THE MEASURE AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND.

**2018 Registry Flow for Quality ID #205 NQF #0409: HIV/AIDS:
Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis**



SAMPLE CALCULATIONS:

Data Completeness=

$$\frac{\text{Performance Met (a=40 patients)} + \text{Denominator Exception (b=20 patients)} + \text{Performance not Met (c=10 patients)}}{\text{Eligible Population / Denominator (d=80 patients)}} = \frac{70 \text{ patients}}{80 \text{ patients}} = 87.50\%$$

Performance Rate=

$$\frac{\text{Performance Met (a= 40 patients)}}{\text{Data Completeness Numerator (70 patients) – Denominator Exception (b=20 patients)}} = \frac{40 \text{ patients}}{50 \text{ patients}} = 80.00\%$$

*See the posted Measure Specification for specific coding and instructions to submit this measure.

NOTE: Submission Frequency: Patient-process

CPT only copyright 2017 American Medical Association. All rights reserved. The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification. v2

**2018 Registry Flow For Quality ID
#205 NQF #0409: HIV/AIDS: Sexually Transmitted Disease Screening for
Chlamydia, Gonorrhea, and Syphilis**

Please refer to the specific section of the specification to identify the denominator and numerator information for use in submitting this Individual Specification. This flow is for registry data submission.

1. Start with Denominator
2. Check Patient Age:
 - a. If the Age is greater than or equal to 13 years of age at Date of Service and equals No during the measurement period, do not include in Eligible Patient Population. Stop Processing.
 - b. If the Age is greater than or equal to 13 years of age at Date of Service and equals Yes during the measurement period, proceed to Check Patient Diagnosis.
3. Check Patient Diagnosis:
 - a. If Diagnosis for HIV/AIDS as Listed In the Denominator equals No, do not include in Eligible Patient Population. Stop Processing.
 - b. If Diagnosis for HIV/AIDS as Listed in the Denominator equals Yes, proceed to Check Encounter Performed.
4. Check Encounter Performed:
 - a. If Encounter as Listed in the Denominator equals No, do not include in Eligible Patient Population. Stop Processing.
 - b. If Encounter as Listed in the Denominator equals Yes, proceed to Check At Least Two Denominator Eligible Encounters During the Measurement Year, With at Least 90 Days Between Each.
5. Check At Least Two Denominator Eligible Encounters During the Measurement Year, With at Least 90 Days Between Each:
 - a. If at Least Two Denominator Eligible Encounters During the Measurement Year, With at Least 90 Days Between Each equals No, do not include in Eligible Patient Population. Stop Processing.
 - b. If at Least Two Denominator Eligible Encounters During the Measurement Year, With at Least 90 Days Between Each equals Yes, proceed to Check Patients Who Use Hospice Services Any Time During the Measurement Period.
6. Check Patients Who Use Hospice Services Any Time During the Measurement Period:
 - a. If Patients Who Use Hospice Services Any Time During the Measurement Period equals No, include in the Eligible Population.
 - b. If Patients Who Use Hospice Services Any Time During the Measurement Period equals Yes, do not include in Eligible Patient Population. Stop Processing.
7. Denominator Population:
 - a. Denominator population is all Eligible Patients in the denominator. Denominator is represented as Denominator in the Sample Calculation listed at the end of this document. Letter d equals 80 patients in the Sample Calculation.
8. Start Numerator
9. Check Chlamydia, Gonorrhea and Syphilis Screenings Documented as Performed:
 - a. If Chlamydia, Gonorrhea and Syphilis Screening Results Documented equals Yes, include in Data Completeness Met and Performance Met.

- b. Data Completeness Met and Performance Met letter is represented as Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a equals 40 patients in Sample Calculation.
 - c. If Chlamydia, Gonorrhea and Syphilis Screening Results Documented as Performed equals No, proceed to Chlamydia, Gonorrhea and Syphilis Screening Results Not Documented, Due to Patient Refusal.
10. Check Chlamydia, Gonorrhea and Syphilis Screening Results Not Documented, Due to Patient Refusal:
- a. If Chlamydia, Gonorrhea and Syphilis Screening Results Not Documented, Due to Patient Refusal equals Yes, include in Data Completeness Met and Denominator Exception.
 - b. Data Completeness Met and Denominator Exception letter is represented as Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter b equals 20 patients in the Sample Calculation.
 - c. If Chlamydia, Gonorrhea and Syphilis Screening Results Not Documented, Due to Patient Refusal equals No, proceed to Check Chlamydia, Gonorrhea and Syphilis Screening Results Not Documented as Performed, Reason Not Given.
11. Check Chlamydia, Gonorrhea and Syphilis Screening Results Not Documented as Performed, Reason Not Given:
- a. If Chlamydia, Gonorrhea and Syphilis Screening Results Not Documented as Performed, Reason Not Given equals Yes, include in Data Completeness Met and Performance Not Met.
 - b. Data Completeness Met and Performance Not Met letter is represented as Data Completeness in the Sample Calculation listed at the end of this document. Letter c equals 10 patients in the Sample Calculation.
 - c. If Chlamydia, Gonorrhea and Syphilis Screening Results Not Documented as Performed, Reason Not Given equals No, proceed to Data Completeness not Met
12. Check Data Completeness Not Met:
- a. If Data Completeness Not Met equals No, Quality Data Code or equivalent not submitted. 10 patients have been subtracted from the Data Completeness Numerator in the Sample Calculation.

SAMPLE CALCULATIONS:

Data Completeness=

$$\frac{\text{Performance Met (a=40 patients)} + \text{Denominator Exception (b=20 patients)} + \text{Performance not Met (c=10 patients)}}{\text{Eligible Population / Denominator (d=80 patients)}} = \frac{70 \text{ patients}}{80 \text{ patients}} = 87.50\%$$

Performance Rate=

$$\frac{\text{Performance Met (a= 40 patients)}}{\text{Data Completeness Numerator (70 patients) – Denominator Exception (b=20 patients) = 50 patients}} = \frac{40 \text{ patients}}{50 \text{ patients}} = 80.00\%$$