

**Quality ID #435: Quality of Life Assessment For Patients With Primary Headache Disorders**  
– National Quality Strategy Domain: Effective Clinical Care  
– Meaningful Measure Area: Patient Reported Functional Outcomes

**2019 COLLECTION TYPE:**  
**MIPS CLINICAL QUALITY MEASURES (CQMS)**

**MEASURE TYPE:**  
Patient Reported Outcome – High Priority

**DESCRIPTION:**  
Percentage of patients with a diagnosis of primary headache disorder whose health related quality of life (HRQoL) was assessed with a tool(s) during at least two visits during the 12 month measurement period AND whose health related quality of life score stayed the same or improved

**INSTRUCTIONS:**  
This measure is to be submitted at least **once per performance period** for patients with a diagnosis of primary headache during the performance period. This measure may be submitted by Merit-based Incentive Payment System (MIPS) eligible clinicians who perform the quality actions described in the measure based on the services provided and the measure-specific denominator coding.

**Measure Submission Type:**  
Measure data may be submitted by individual MIPS eligible clinicians, groups, or third party intermediaries. The listed denominator criteria are used to identify the intended patient population. The numerator options included in this specification are used to submit the quality actions as allowed by the measure. The quality-data codes listed do not need to be submitted by MIPS eligible clinicians, groups, or third party intermediaries that utilize this modality for submissions; however, these codes may be submitted for those third party intermediaries that utilize Medicare Part B claims data. For more information regarding Application Programming Interface (API), please refer to the Quality Payment Program (QPP) website.

**DENOMINATOR:**  
All patients with a diagnosis of primary headache disorder

- Denominator Instruction:**  
Primary Headache for the purpose of this measure includes the following types of headache:
- **Migraine:** Migraine without aura, migraine with aura, childhood periodic syndromes that are commonly precursors of migraine, retinal migraine, complications of migraine, probable migraine
  - **Tension-Type Headache (TTH):** Infrequent episodic TTH, frequent episodic TTH, chronic TTH, probable TTH.
  - **Cluster Headache (CH) and Other Trigeminal Autonomic Cephalgias:** Cluster headache, paroxysmal hemicrania, short-lasting unilateral neuralgia form headache attacks with conjunctival injection and tearing (SUNCT), probable trigeminal autonomic cephalgia
  - **Other Primary Headaches:** Primary stabbing headache, primary cough headache, primary exertional headache, primary headache associated with sexual activity, hypnic headache, primary thunderclap headache, hemicrania continua, new daily-persistent headache.

***DENOMINATOR NOTE:*** \*Signifies that this CPT Category I code is a non-covered service under the Medicare Part B Physician Fee Schedule (PFS). These non-covered services should be counted in the denominator population for MIPS CQMs.

**Denominator Criteria (Eligible Cases):**

**Diagnosis for Primary Headache (ICD-10-CM):** G43.001, G43.009, G43.011, G43.019, G43.101, G43.109, G43.111, G43.119, G43.401, G43.409, G43.411, G43.419, G43.501, G43.509, G43.511, G43.519, G43.601, G43.609, G43.611, G43.619, G43.701, G43.709, G43.711, G43.719, G43.801, G43.809, G43.811, G43.819, G43.821, G43.829, G43.831, G43.839, G43.901, G43.909, G43.911, G43.919, G44.009, G44.019, G44.029, G44.039, G44.049, G44.059, G44.099, G44.1, G44.209, G44.219, G44.221, G44.229, G44.51, G44.52, G44.53, G44.59, G44.81, G44.82, G44.83, G44.84, G44.85, G44.89

**AND**

**Encounter during the performance period (CPT):** 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241\*, 99242\*, 99243\*, 99244\*, 99245\*

**WITHOUT**

**Telehealth Modifier:** GQ, GT, 95, POS 02

**AND**

**At Least Two Visits during the Performance Period**

**NUMERATOR:**

Patient whose health related quality of life was assessed with a tool(s) during at least two visits during the 12 month measurement period AND whose health related quality of life score stayed the same or improved

**Numerator Instructions:**

**List quality of life (QoL) tools:** Migraine Disability Assessment (MIDAS) and PedMIDAS(proprietary); Headache Impact Test-6 (HIT-6)(proprietary); Migraine Specific Quality of Life Tool (MSQ); Neck Disability Index (NDI)-used for cervicogenic headaches; McGill Questionnaire.

**Timing Between Visits:** Must be separated by at least 90 days for MIDAS and at least 4 weeks for any other tool.

**See specific tools for scoring methods related to improvement or stayed the same:** Each tool defines improvement differently based on their scoring methodology. For example, when using the MIDAS improvement would be indicated by reduction in MIDAS disability grade and in the HIT-6 a reduction in the number of days with disability overtime indicates improvement.

**NUMERATOR NOTE:** *The two assessments must be separated by at least 90 days for MIDAS and at least 4 weeks for any other tool. It is expected that the QoL score or ranking will stay the same or improve in order for this measure to be successfully completed. Denominator Exception(s) are determined on the date of the denominator eligible encounter.*

**Numerator Options:**

**Performance Met:**

Health-related quality of life assessed with tool during at least two visits and quality of life score remained the same or improved (**G9634**)

**OR**

**Denominator Exception:**

Health-related quality of life not assessed with tool for documented reason(s) (e.g., patient has a cognitive or neuropsychiatric impairment that impairs his/her ability to complete the HRQoL survey, patient has the inability to read and/or write in order to complete the HRQoL questionnaire) (**G9635**)

**OR**

**Performance Not Met:**

Health-related quality of life not assessed with tool during at least two visits or quality of life score declined (**G9636**)

## **RATIONALE:**

This measure establishes an initial or baseline QoL score from which the patient should use the same QoL tool/questionnaire at least one additional time during the measurement period. The two assessments must be separated by at least 90 days for MIDAS and at least 4 weeks for any other tool.<sup>3</sup> It is expected that the QoL score or ranking will stay the same or improve in order for this measure to be successfully completed.

### Gap in Care

Migraine impacts a person's function in different activity domains during attacks. HRQoL is affected both during and after attacks.<sup>1</sup> Migraine reduces HRQoL more than osteoarthritis or diabetes.<sup>2</sup> In the US and UK, subjects with migraine had lower scores ( $p < 0.001$ ) on both the Mental Component Score (MCS-12) and Physical Component Score (PCS-12) than their non-migraine counterparts. Significant differences were maintained after controlling for gender, age, and education. Migraine and depression were highly correlated (adjusted prevalence ratio 2.7, 95% CI

2.1 to 3.5). Further, migraine and depression are highly associated with attack frequency (for MCS-12 and PCS-12) and disability (MCS-12). Subjects with migraine selected from the general population have lower HRQoL as measured by the Short Form (SF-morbid) and each exerts a significant and independent influence on HRQoL.<sup>3</sup>

### Opportunity for Improvement

This is the first clinician level patient reported outcome measure (PROM) focused on maintaining or improving the QoL of patients with primary headache disorders. The Work Group felt that even though the majority evidence is focused on migraine that patients with other primary headache disorders could greatly benefit from the utilization of this measure.

The use of PROMs to investigate levels of disability and HRQoL are increasingly being used in headache services research. HRQoL and disability are positively impacted by treatment interventions.<sup>4</sup> Health care professionals often do not recognize the degree and the scope of functional impairment imposed by migraines. There is a missed opportunity for clinicians to effectively communicate with the patient to understand their headache-related disability and appropriately prescribe acute, prophylactic, or biobehavioral treatments. This measure has the potential to reduce personal and societal costs of headache disorders offering a continuity of care.

1Buse DC, Rupnow MF, Lipton RB. Assessing and managing all aspects of migraine: migraine attacks, migraine related functional impairment, common comorbidities, and quality of life. *Mayo Clin Proc* 2009; 84: 422-435

2Buse DC, Manack AN, Fanning KM, et al. Chronic Migraine Prevalence, Disability, and Sociodemographic Factors: Results From the American Migraine Prevalence and Prevention Study. *Headache*. 2012 Jun 22. doi: 10.1111/j.1526-4610.2012.02223.x. [Epub ahead of print]

3Lipton RB, Hamelsky SW, Kolodner KB et al. Migraine, quality of life, and depression A population-based case-control study *Neurology*, 2000 vol. 55 no. 5 629-635

4D'Amico D, Grazi L, Usai S, Leonardi M. Disability and quality of life in headache: where are we not and where we are heading. *Neurol Sci* 2013 34(S1):S1-S5

## **CLINICAL RECOMMENDATION STATEMENTS:**

Discuss the benefits and risks of prophylactic treatment for migraine with the person, taking into account the person's preference, comorbidities, risk of adverse events and the impact of the headache on their QoL. (No level of evidence)

1

Compared with people without headache and to people with other chronic conditions, people with headache report compromised physical, mental, and social functioning, particularly those with a high frequency of attack. People with

headache reported diminished functioning and well-being on all eight domains as compared with people without headache .2

1. NICE Headaches: Diagnosis and management of headaches in young people and adults. National Clinical Guideline Centre on behalf of the National Institute for Health and Clinical Excellence (NICE) September 2012; NICE clinical guideline 150
2. Terwindt GM, Ferrari MD, Tihuis M et al. The impact of migraine on quality of life in the general population: The GEM study Neurology 2000 55:624-629


**COPYRIGHT:**

© 2015 American Academy of Neurology Institute All rights reserved.

Limited proprietary coding is contained in the measure specifications for convenience. Users of the proprietary coding sets should obtain all necessary licenses from the owners of these code sets. The AAN and its members disclaim all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications.

CPT ® is a registered trademark of the American Medical Association.

**2019 Clinical Quality Measure Flow for Quality ID #435:  
Quality of Life Assessment For Patients With Primary Headache Disorders**



**SAMPLE CALCULATIONS:**

**Data Completeness=**  
 Performance Met (a=40 patients) + Denominator Exception (b=10 patients) + Performance Not Met (c=20 patients) = 70 patients = 87.50%  
 Eligible Population / Denominator (d=80 patients) = 80 patients

**Performance Rate=**  
 $\frac{\text{Performance Met (a=40 patients)}}{\text{Data Completeness Numerator (70 patients) - Denominator Exception (b=10 patients)}} = \frac{40 \text{ patients}}{60 \text{ patients}} = 66.67\%$

\*See the posted Measure Specification for specific coding and instructions to submit this measure.

NOTE : Submission Reporting Frequency: Patient-Intermediate

CPT only copyright 2018 American Medical Association. All rights reserved.  
 The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.

**2019 Clinical Quality Measure Flow Narrative for Quality ID #435:  
Quality of Life Assessment For Patients With Primary Headache Disorders**

Please refer to the specific section of the specification to identify the denominator and numerator information for use in submitting this Individual Specification.

1. Start with Denominator
2. Check Patient Diagnosis:
  - a. If Diagnosis for Primary Headache as Listed in the Denominator equals No, do not include in Eligible Population. Stop Processing.
  - b. If Diagnosis for Primary Headache as Listed in the Denominator equals Yes, proceed to check Encounter Performed.
3. Check Encounter Performed:
  - a. If Encounter as Listed in the Denominator equals No, do not include in Eligible Population. Stop Processing.
  - b. If Encounter as Listed in the Denominator equals Yes, proceed to check Telehealth Modifier.
4. Check Telehealth Modifier:
  - a. If Telehealth Modifier equals Yes, do not include in Eligible Population. Stop Processing.
  - b. If Telehealth Modifier equals No, proceed to check At Least Two Visits During the Performance Period.
5. Check At Least Two Visits During the Performance Period:
  - a. If At Least Two Visits During the Performance Period equals No, do not include in Eligible Population. Stop Processing.
  - b. If At Least Two Visits During the Performance Period equals Yes, include in Eligible Population.
6. Denominator Population:
  - a. Denominator Population is all Eligible Patients in the Denominator. Denominator is represented as Denominator in the Sample Calculation listed at the end of this document. Letter d equals 80 patients in the Sample Calculation.
7. Start Numerator
8. Check Health-Related Quality of Life Assessed with Tool During at Least Two Visits and Quality of Life Score Remained the Same or Improved:
  - a. If Health-Related Quality of Life Assessed with Tool During at Least Two Visits and Quality of Life Score Remained the Same or Improved equals Yes, include in Data Completeness Met and Performance Met.
  - b. Data Completeness Met and Performance Met letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a equals 40 patients in the Sample Calculation.

- c. If Health-Related Quality of Life Assessed with Tool During at Least Two Visits and Quality of Life Score Remained the Same or Improved equals No, proceed to check Health-Related Quality of Life Not Assessed with Tool for Documented Reason(s).
9. Check Health-Related Quality of Life Not Assessed with Tool for Documented Reason(s):
- a. If Health-Related Quality of Life Not Assessed with Tool for Documented Reason(s) equals Yes, include in Data Completeness Met and Denominator Exception.
  - b. Data Completeness Met and Denominator Exception letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter b equals 10 patients in the Sample Calculation.
  - c. If Health-Related Quality of Life Not Assessed with Tool for Documented Reason(s) equals No, proceed to check Health-Related Quality of Life Not Assessed with Tool During at Least Two Visits or Quality of Life Score Declined.
10. Check Health-Related Quality of Life Not Assessed with Tool During at Least Two Visits or Quality of Life Score Declined:
- a. If Health-Related Quality of Life Not Assessed with Tool During at Least Two Visits or Quality of Life Score Declined equals Yes, include in Data Completeness Met and Performance Not Met.
  - b. Data Completeness Met and Performance Not Met letter is represented in the Data Completeness in the Sample Calculation listed at the end of this document. Letter c equals 20 patients in the Sample Calculation.
  - c. If Health-Related Quality of Life Not Assessed with Tool During at Least Two Visits or Quality of Life Score Declined equals No, proceed to check Data Completeness Not Met
11. Check Data Completeness Not Met:
- a. If Data Completeness Not Met, the Quality Data Code or equivalent was not submitted. 10 patients have been subtracted from the Data Completeness Numerator in the Sample Calculation.

**SAMPLE CALCULATIONS:**

**Data Completeness=**

$$\frac{\text{Performance Met (a=40 patients)} + \text{Denominator Exception (b=10 patients)} + \text{Performance Not Met (c=20 patients)}}{\text{Eligible Population / Denominator (d=80 patients)}} = \frac{70 \text{ patients}}{80 \text{ patients}} = 87.50\%$$

**Performance Rate=**

$$\frac{\text{Performance Met (a=40 patients)}}{\text{Data Completeness Numerator (70 patients) - Denominator Exception (b=10 patients)}} = \frac{40 \text{ patients}}{60 \text{ patients}} = 66.67\%$$