

**Quality ID #451 (NQF 1859): RAS (KRAS and NRAS) Gene Mutation Testing Performed for Patients with Metastatic Colorectal Cancer who Receive Anti-epidermal Growth Factor Receptor (EGFR) Monoclonal Antibody Therapy**  
– National Quality Strategy Domain: Effective Clinical Care  
– Meaningful Measure Area: Appropriate Use of Healthcare

**2021 COLLECTION TYPE:**  
**MIPS CLINICAL QUALITY MEASURES (CQMS)**

**MEASURE TYPE:**  
Process

**DESCRIPTION:**  
Percentage of adult patients (aged 18 or over) with metastatic colorectal cancer who receive anti-epidermal growth factor receptor monoclonal antibody therapy for whom RAS (KRAS and NRAS) gene mutation testing was performed

**INSTRUCTIONS:**  
This measure is to be submitted **once per performance period** for patients with colorectal cancer seen during the performance period. This measure may be submitted by Merit-based Incentive Payment System (MIPS) eligible clinicians who perform the quality actions described in the measure based on the services provided and the measure-specific denominator coding.

**Measure Submission Type:**  
Measure data may be submitted by individual MIPS eligible clinicians, groups, or third-party intermediaries. The listed denominator criteria are used to identify the intended patient population. The numerator options included in this specification are used to submit the quality actions as allowed by the measure. The quality-data codes listed do not need to be submitted by MIPS eligible clinicians, groups, or third-party intermediaries that utilize this modality for submissions; however, these codes may be submitted for those third-party intermediaries that utilize Medicare Part B claims data. For more information regarding Application Programming Interface (API), please refer to the Quality Payment Program (QPP) website.

**DENOMINATOR:**  
Adult patients with metastatic colorectal cancer who receive anti-EGFR monoclonal antibody therapy

**Denominator Criteria (Eligible Cases):**

Patients aged ≥ 18 years on date of encounter

**AND**

**Diagnosis of initial colon or rectal cancer (ICD-10 CM):** C18.0, C18.2, C18.3, C18.4, C18.5, C18.6, C18.7, C18.8, C18.9, C19, C20

**AND**

**Patient Encounter during the performance period (CPT):** 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215

**WITHOUT**

**Telehealth Modifier:** GQ, GT, 95, POS 02

**AND**

**Two or more encounters at the reporting site**

**AND**

**Patient has metastatic disease at diagnosis:** G9838

**AND**

**Anti-EGFR monoclonal antibody therapy:** G9839

## **NUMERATOR:**

RAS (KRAS and NRAS) gene mutation testing performed before initiation of anti-EGFR MoAb

### **Definition:**

**RAS mutation testing** - RAS testing for this measure refers to assays that detect mutations in codons 12 and 13 of exon 2, codons 59 and 61 of exon 3 and codons 117 and 146 in exon 4 in KRAS or NRAS. Do not include results from mutations at other codons or assays for other alterations (e.g., BRAF, PI3K, PTEN genes). The College of American Pathologists (CAP) Perspectives on Emerging Technology (POET) Report on RAS mutation testing provides additional guidance on testing.

If multiple RAS mutation tests have been performed, refer to the most recent test results.

Anti-EGFR monoclonal antibody includes cetuximab or panitumumab.

### **Numerator Instructions:**

In the absence of any documentation regarding testing for the RAS (KRAS and NRAS) gene mutation, submit **G9841**: RAS (KRAS and NRAS) gene mutation testing not performed before initiation of anti-EGFR MoAb. Report **G9840**: RAS (KRAS and NRAS) gene mutation testing performed before initiation of anti-EGFR MoAb, if the report indicates a mutation within codons 12 and 13 of exon 2, codons 59 and 61 of exon 3 and codons 117 and 146 in exon 4 in KRAS or NRAS, where KRAS or NRAS gene was detected in the DNA extracted from the colon tumor specimen.

### **Numerator Options:**

#### ***Performance Met:***

RAS (KRAS and NRAS) gene mutation testing performed before initiation of anti-EGFR MoAb (**G9840**)

## **OR**

#### ***Performance Not Met:***

RAS (KRAS and NRAS) gene mutation testing not performed before initiation of anti-EGFR MoAb (**G9841**)

## **RATIONALE:**

The American Society of Clinical Oncology (ASCO) envisions that use of this measure will improve concordance with recommendations for RAS (KRAS and NRAS) testing for patients with metastatic colorectal cancer. We recognize the importance of ensuring that the appropriate patient population receives guideline concordant treatment as studies demonstrate that the administration of EGFR-targeted therapies, specifically cetuximab or panitumumab, offer no clinical benefit to patients diagnosed with KRAS-mutated or NRAS-mutated tumors. Clinical trial data strongly suggest that patients with RAS mutations are better served with other targeted therapies, especially considering the harms and costs of anti-EGFR treatment. Therefore, the measure focus is on halting use of anti-EGFR MoAb therapies in patients who will not derive any benefit.

## **CLINICAL RECOMMENDATION STATEMENTS:**

This measure is based on ASCO and NCCN Guidelines:

“Colorectal carcinoma patients being considered for anti-EGFR therapy must receive RAS mutational testing. Mutational analysis should include KRAS and NRAS codons 12, 13 of exon 2; 59, 61 of exon 3; and 117 and 146 of exon 4 (“expanded” or “extended” RAS)”.

Sepulveda AR, Hamilton SR, Allegra CJ, et al: Molecular Biomarkers for the Evaluation of Colorectal Cancer: Guideline From the American Society for Clinical Pathology, College of American Pathologists, Association for Molecular Pathology, and the American Society of Clinical Oncology. Journal of Clinical Oncology 35:1453-1486, 2017.

“All patients with metastatic colorectal cancer should have tumor tissue genotyped for RAS (KRAS and NRAS) and BRAF mutations individually or as part of a next-generation sequencing (NGS) panel. Patients with any known KRAS mutation (exon 2, 3, 4) or NRAS mutation (exon 2, 3, 4) should not be treated with either cetuximab or panitumumab.”

“A sizeable body of literature has shown that tumors with a mutation in codon 12 or 13 of exon 2 of the KRAS gene are essentially insensitive to cetuximab or panitumumab therapy... More recent evidence shows mutations in KRAS outside of exon 2 and mutations in NRAS are also predictive for a lack of benefit of anti-EGFR therapies.

The panel therefore strongly recommends RAS (KRAS/NRAS) genotyping of tumor tissue (either primary tumor or metastasis) in all patients with metastatic colorectal cancer. Patients with known KRAS or NRAS mutations should not be treated with either cetuximab or panitumumab, either alone or in combination with other anticancer agents, because they have virtually no chance of benefit and the exposure to toxicity and expense cannot be justified. It is implied throughout the guidelines that NCCN recommendations involving cetuximab or panitumumab relate only to patients with disease characterized by RAS wild-type genes. ASCO released a Provisional Clinical Opinion Update on extended RAS testing in patients with metastatic colorectal cancer that is consistent with the NCCN Panel's recommendations. A guideline on molecular biomarkers for colorectal cancer developed by the ASCP, CAP, AMP and ASCO also recommends RAS testing consistent with the NCCN recommendations” (MS-43)

NCCN Clinical Practice Guidelines in Oncology™. Colon Cancer, V.3.2019 (MS-30)  
[https://www.nccn.org/professionals/physician\\_gls/pdf/colon.pdf](https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf)

### **COPYRIGHT:**

The Measure is not clinical guidelines, does not establish a standard of medical care, and has not been tested for all potential applications.

The Measure, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, e.g., use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the Measures for commercial gain, or incorporation of the Measure into a product or service that is sold, licensed or distributed for commercial gain.

Commercial uses of the Measure require a license agreement between the user and the American Society of Clinical Oncology (ASCO) and prior written approval of ASCO. Neither ASCO nor its members shall be responsible for any use of the Measures.

ASCO encourages use of the Measures by other health care professionals, where appropriate.

### **THE MEASURES AND SPECIFICATIONS ARE PROVIDED “AS IS” WITHOUT WARRANTY OF ANY KIND.**


© 2021 American Society of Clinical Oncology. All Rights Reserved.

Limited proprietary coding is contained in the Measure specification for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets. ASCO and its members disclaim all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specification.

CPT® contained in the Measures specifications is copyright 2004-2020 American Medical Association. ICD-10 is copyright 2020 World Health Organization. LOINC® copyright 2004-2020 Regenstrief Institute, Inc. SNOMED CLINICAL TERMS (SNOMED CT®) copyright 2004-2020 College of American Pathologists. All Rights Reserved.

**2021 Clinical Quality Measure Flow for Quality ID #451 (NQF 1859):  
RAS (KRAS and NRAS) Gene Mutation Testing Performed for Patients with Metastatic Colorectal  
Cancer who Receive Anti-epidermal Growth Factor Receptor (EGFR) Monoclonal Antibody Therapy**

*Disclaimer: Refer to the measure specification for specific coding and instructions to submit this measure.*



### **SAMPLE CALCULATIONS**

**Data Completeness=**

$$\frac{\text{Performance Met (a=40 patients)} + \text{Performance Not Met (c=30 patients)}}{\text{Eligible Population / Denominator (d=80 patients)}} = \frac{70 \text{ patients}}{80 \text{ patients}} = 87.50\%$$

**Performance Rate=**

$$\frac{\text{Performance Met (a=40 patients)}}{\text{Data Completeness Numerator (70 patients)}} = \frac{40 \text{ patients}}{70 \text{ patients}} = 57.14\%$$

\*See the posted measure specification for specific coding and instructions to submit this measure.

NOTE: Submission Frequency: Patient-Process

CPT only copyright 2020 American Medical Association. All rights reserved.  
The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.

v5

**2020 Clinical Quality Measure Flow Narrative for Quality ID #451 NQF #1859:  
RAS (KRAS and NRAS) Gene Mutation Testing Performed for Patients with Metastatic Colorectal Cancer  
who Receive Anti-epidermal Growth Factor Receptor (EGFR) Monoclonal Antibody Therapy**

*Disclaimer: Refer to the measure specification for specific coding and instructions to submit this measure.*

1. Start with Denominator
2. Check *Patients aged is greater than or equal to 18 years on date of encounter*:
  - a. If *Patients aged is greater than or equal to 18 years on date of encounter* equals No, do not include in *Eligible Population/Denominator*. Stop Processing.
  - b. If *Patients age is greater than or equal to 18 years on date of encounter* equals Yes, proceed to check *Diagnosis of initial colon or rectal cancer as listed in Denominator\**.
3. Check *Diagnosis of initial colon or rectal cancer as listed in Denominator\**:
  - a. If *Diagnosis of initial colon or rectal cancer as listed in Denominator\** equals No, do not include in *Eligible Population/Denominator*. Stop Processing.
  - b. If *Diagnosis of initial colon or rectal cancer as listed in Denominator\** equals Yes, proceed to check *Patient encounter during the performance period as listed in Denominator\**.
4. Check *Patient encounter during the performance period as listed in Denominator\**:
  - a. If *Patient encounter during the performance period as listed in Denominator\** equals No, do not include in *Eligible Population/Denominator*. Stop Processing.
  - b. If *Patient encounter during the performance period as listed in Denominator\** equals Yes, proceed to check *Telehealth Modifier*.
5. Check *Telehealth Modifier*:
  - a. If *Telehealth Modifier* equals No, do not include in *Eligible Population/Denominator*. Stop Processing.
  - b. If *Telehealth Modifier* equals Yes, proceed to check *Two or more encounters at reporting site*.
6. Check *Two or more encounters at reporting site*:
  - a. If *Two or more encounters at reporting site* equals No, do not include in *Eligible Population/Denominator*. Stop Processing.
  - b. *Two or more encounters at reporting site* equals Yes, proceed to check *Patient has metastatic disease at diagnosis*.
7. Check *Patient has metastatic disease at diagnosis*:
  - a. If *Patient has metastatic disease at diagnosis* equals No, do not include in *Eligible Population/Denominator*. Stop Processing.
  - b. If *Patient has metastatic disease at diagnosis* equals Yes, proceed to check *Anti-EGFR monoclonal antibody therapy*.
8. Check *Anti-EGFR monoclonal antibody therapy*:

- a. If *Anti-EGFR monoclonal antibody therapy* equals No, do not include in *Eligible Population/Denominator*. Stop Processing.
  - b. If *Anti-EGFR monoclonal antibody therapy* equals Yes, include in *Eligible Population/Denominator*.
9. Denominator Population:
- a. Denominator Population is all Eligible Patients in the Denominator. Denominator is represented as Denominator in the Sample Calculation listed at the end of this document. Letter d equals 80 patients in the Sample Calculation.
10. Start Numerator
11. Check *RAS (KRAS and NRAS) gene mutation testing performed before initiation of anti-EGFR MoAb*:
- a. If *RAS (KRAS and NRAS) gene mutation testing performed before initiation of anti-EGFR MoAb* equals Yes, include in *Data Completeness Met and Performance Met*.
    - *Data Completeness Met and Performance Met* letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a equals 40 patients in Sample Calculation.
  - b. If *RAS (KRAS and NRAS) gene mutation testing performed before initiation of anti-EGFR MoAb* equals No, proceed to check *RAS (KRAS and NRAS) gene mutation testing not performed before initiation of anti-EGFR MoAb*.
12. Check *RAS (KRAS and NRAS) gene mutation testing not performed before initiation of anti-EGFR MoAb*:
- a. If *RAS (KRAS and NRAS) gene mutation testing not performed before initiation of anti-EGFR MoAb* equals Yes, include in *Data Completeness Met and Performance Not Met*.
    - *Data Completeness Met and Performance Not Met* letter is represented in the Data Completeness in the Sample Calculation listed at the end of this document. Letter c equals 30 patients in the Sample Calculation.
  - b. If *RAS (KRAS and NRAS) gene mutation testing not performed before initiation of anti-EGFR MoAb* equals No, proceed to check *Data Completeness Not Met*.
13. Check *Data Completeness Not Met*:
- a. If *Data Completeness Not Met*, the Quality Data Code or equivalent was not submitted. 10 patients have been subtracted from the Data Completeness Numerator in the Sample Calculation.

**Sample Calculations:**

Data Completeness equals Performance Met (a equals 40 patients) plus Performance Not Met (c equals 30 patients) divided by Eligible Population / Denominator (d equals 80 patients). All equals 70 patients divided by 80 patients. All equals 87.50 percent.

Performance Rate equals Performance Met (a equals 40 patients) divided by Data Completeness Numerator (70 patients). All equals 40 patients divided by 70 patients. All equals 57.14 percent.

\*See the posted measure specification for specific coding and instructions to submit this measure.

NOTE: Submission Frequency: Patient-Process

The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.