Quality ID #185: Colonoscopy Interval for Patients with a History of Adenomatous Polyps – Avoidance of Inappropriate Use

- National Quality Strategy Domain: Communication and Care Coordination

- Meaningful Measure Area: Appropriate Use of Healthcare

#### 2021 COLLECTION TYPE: MIPS CLINICAL QUALITY MEASURES (CQMS)

## MEASURE TYPE:

Process – High Priority

### **DESCRIPTION:**

Percentage of patients aged 18 years and older receiving a surveillance colonoscopy, with a history of prior adenomatous polyp(s) in previous colonoscopy findings, which had an interval of 3 or more years since their last colonoscopy

### **INSTRUCTIONS:**

This measure is to be submitted **<u>each time</u>** a surveillance colonoscopy is performed during the performance period. It is anticipated the Merit-based Incentive Payment System (MIPS) eligible clinician who performs the listed procedures, as specified in the denominator coding, will submit on this measure. Patients who have a coded colonoscopy procedure that has a modifier 52, 53, 73 or 74 will not qualify for inclusion into this measure.

### Measure Submission Type:

Measure data may be submitted by individual MIPS eligible clinicians, groups, or third party intermediaries. The listed denominator criteria are used to identify the intended patient population. The numerator options included in this specification are used to submit the quality actions as allowed by the measure. The quality-data codes listed do not need to be submitted by MIPS eligible clinicians, groups, or third party intermediaries that utilize this modality for submissions; however, these codes may be submitted for those third party intermediaries s that utilize Medicare Part B claims data. For more information regarding Application Programming Interface (API), please refer to the Quality Payment Program (QPP) website.

# **DENOMINATOR:**

All patients aged 18 years and older receiving a surveillance colonoscopy, with a history of a prior adenomatous polyp(s) in previous colonoscopy findings

# **Denominator Instructions:**

**MIPS** eligible clinicians who indicate that the colonoscopy procedure is incomplete or was discontinued should use the procedure number and the addition (as appropriate) of modifier 52, 53, 73, or 74. Patients who have a coded colonoscopy procedure that has a modifier 52, 53, 73, or 74 will not qualify for inclusion into this measure.

#### Denominator Criteria (Eligible Cases):

Patients aged ≥ 18 years on date of encounter <u>AND</u> Diagnosis for history of adenomatous (colonic) polyp(s) (ICD-10-CM): Z86.010 <u>AND</u> Patient procedure during the performance period (CPT or HCPCS): 44388, 44389, 44392, 44394, 45378, 45380, 45381, 45384, 45385, G0105 <u>WITHOUT</u> Modifiers: 52, 53, 73 or 74

#### NUMERATOR:

Patients who had an interval of 3 or more years since their last colonoscopy

<u>OR</u>	Numerator Options: Performance Met:	Interval of 3 or more years since patient's last colonoscopy, documented <b>(0529F)</b>
	Denominator Exception:	Documentation of medical reason(s) for an interval of less than 3 years since the last colonoscopy (e.g.,last colonoscopy incomplete, last colonoscopy had inadequate prep, piecemeal removal of adenomas, last colonoscopy found greater than 10 adenomas, or patient at high risk for colon cancer [Crohn's disease, ulcerative colitis, lower gastrointestinal bleeding, personal or family history of colon cancer]) <b>(0529F</b> <i>with</i> <b>1P)</b>
	<u>OR</u>	
<u>OR</u>	Denominator Exception:	Documentation of system reason(s) for an interval of less than 3 years since the last colonoscopy (e.g., unable to locate previous colonoscopy report, previous colonoscopy report was incomplete) (0529F with 3P)
	Performance Not Met:	Interval of less than 3 years since patient's last colonoscopy, reason not otherwise specified <b>(0529F <i>with</i> 8P)</b>

#### RATIONALE:

Colorectal cancer is the 2nd leading cause of cancer death in the United States. Colonoscopy is the recommended method of surveillance after the removal of adenomatous polyps because it has been shown to significantly reduce subsequent colorectal cancer incidence. The time interval for the development of malignant changes in adenomatous polyps is estimated at 5 to 25 years. (ICSI, 2006) Inappropriate interval recommendations can result in overuse of resources and can lead to significant patient harm. Performing colonoscopy too often not only increases patients' exposure to procedural harm, but also drains resources that could be more effectively used to adequately screen those in need. (Lieberman et al, 2009)

#### **CLINICAL RECOMMENDATION STATEMENTS:**

Patients with only 1 or 2 small (< 1 cm) tubular adenomas with only low-grade dysplasia should have their next followup colonoscopy in 5–10 years; the precise timing within this interval should be based on other clinical factors (such as prior colonoscopy findings, family history, and the preferences of the patient and judgment of the physician). Patients with 3 to 10 adenomas, or any adenoma  $\geq$  1 cm, or any adenoma with villous features, or high-grade dysplasia should have their next follow-up colonoscopy in 3 years providing that piecemeal removal has not been performed and the adenoma(s) are removed completely; if the follow-up colonoscopy is normal or shows only 1 or 2 small tubular adenomas with low-grade dysplasia, then the interval for the subsequent examination should be 5 years. (Winawer, et al, 2006)

Patients with > 10 adenomas are thought to be at particularly high risk, and current multi-society guidelines therefore recommend early surveillance colonoscopy in these individuals (< 3 years). (Lieberman, et al, 2012) However, it is important to note that risk is a continuum; an individual with 11 adenomas is not at dramatically higher risk than an individual with 9 or 10 adenomas. Thus, the optimal threshold at which early surveillance colonoscopy becomes worthwhile is subject to debate. For instance, in the United Kingdom, early surveillance colonoscopy is recommended for individuals with even fewer adenomas ( $\geq$  5 adenomas of any size, or  $\geq$  3 adenomas with at least one large adenoma). A lower threshold is likely to result in higher colonoscopy utilization, but it may also provide greater clinical benefit. (Martinez, et al, 2012)

### **COPYRIGHT**

The Measures are not clinical guidelines, do not establish a standard of medical care, and have not been tested for all potential applications.

The Measures, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, eg, use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the Measures for commercial gain, or incorporation of the Measures into a product or service that is sold, licensed or distributed for commercial gain.

Commercial uses of the Measures require a license agreement between the user and the American Medical Association (AMA), or the American Gastroenterological Association (AGA), or American Society for Gastrointestinal Endoscopy (ASGE) or the American College of Gastroenterology (ACG). Neither the AMA, AGA, ASGE, ACG, nor its members shall be responsible for any use of the Measures.

The AMA's and National Committee for Quality Assurance's significant past efforts and contributions to the development and updating of the Measures is acknowledged. AGA, ASGE and ACG are solely responsible for the review and enhancement ("Maintenance") of the Measures as of August 14, 2014.

AGA, ASGE and ACG encourage use of the Measures by other health care professionals, where appropriate.

#### THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANYKIND.


© 2020 American Medical Association, American Gastroenterological Association, American Society for Gastrointestinal Endoscopy and American College of Gastroenterology. All Rights Reserved. Applicable FARS/DFARS Restrictions Apply to Government Use. For the Merit-Based Incentive Payment System, American Gastroenterological Association is the primary steward for measure revisions.

Limited proprietary coding is contained in the Measure specifications for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets. The AMA, AGA, ASGE, ACG and its members disclaim all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications.

CPT<sup>®</sup> contained in the Measure specifications is copyright 2004-2020 American Medical Association. ICD-10 is copyright 2020 World Health Organization. LOINC<sup>®</sup> copyright 2004-2020 Regenstrief Institute, Inc. This material contains SNOMED CLINICAL TERMS (SNOMED CT<sup>®</sup>) copyright 2004-2020 International Health Terminology Standards Development Organisation. College of American Pathologists. All Rights Reserved.

#### 2021 Clinical Quality Measure Flow for Quality ID #185: Colonoscopy Interval for Patients with a History of Adenomatous Polyps – Avoidance of Inappropriate Use

Disclaimer: Refer to the measure specification for specific coding and instructions to submit this measure.



#### SAMPLE CALCULATIONS

#### Data Completeness=

Data Completeness-			
Performance Met (a=30 procedures) + Denominator Exceptions (b <sup>1</sup> +b <sup>2</sup> =20 procedures) + Performa	ance Not Met (c=20)	procedures) = 70 procedures = 87.50%	
Eligible Population / Denominator (d=80 procedures)		= 80 procedures	
Performance Rate=			
Performance Met (a=30 procedures)	= <u>30 procedures</u>	<u>s</u> = 60.00%	
Data Completeness Numerator (70 procedures) – Denominator Exceptions (b <sup>1</sup> +b <sup>2</sup> =20 procedures)	= 50 procedures	S	

\* See the posted measure specification for specific coding and instructions to submit this measure. NOTE: Submission Frequency: Procedure

CPT only copyright 2020 American Medical Association. All rights reserved. The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification. v5

# 2021 Clinical Quality Measure Flow Narrative for Quality ID #185: Colonoscopy Interval for Patients with a History of Adenomatous Polyps – Avoidance of Inappropriate Use

Disclaimer: Refer to the measure specification for specific coding and instructions to submit this measure.

- 1. Start with Denominator
- 2. Check Patients aged greater than or equal to 18 years on date of encounter.
  - a. If Patients aged greater than or equal to 18 years on date of encounter equals No, do not include in *Eligible Population/Denominator*. Stop processing.
  - b. If Patients aged greater than or equal to 18 years on date of encounter equals Yes, proceed to check Diagnosis for history of adenomatous (colonic) polyp(s) as listed in Denominator\*.
- 3. Check Diagnosis for history of adenomatous (colonic) polyp(s) as listed in Denominator\*:
  - a. If *Diagnosis for history of adenomatous (colonic) polyp(s) as listed in Denominator\** equals No, do not include in *Eligible Population/Denominator*. Stop processing.
  - b. If Diagnosis for history of adenomatous (colonic) polyp(s) as listed in Denominator\* equals Yes, proceed to check Patient procedure during the performance period as listed in Denominator\*.
- 4. Check Patient procedure during the performance period as listed in Denominator\*:
  - a. If *Patient procedure during the performance period as listed in Denominator*\* equals No, do not include in *Eligible Population/Denominator*. Stop processing.
  - b. If Patient procedure during the performance period as listed in Denominator\* equals Yes, proceed to check Modifiers.
- 5. Check Modifiers:
  - a. If Modifiers equals No, include in Eligible Population/Denominator.
  - b. If Modifiers equals Yes, do not include in Eligible Population/Denominator. Stop processing.
- 6. Denominator Population:
  - Denominator Population is all Eligible Procedures in the Denominator. Denominator is represented as Denominator in the Sample Calculation listed at the end of this document. Letter d equals 80 procedures in the Sample Calculation.
- 7. Start Numerator
- 8. Check Interval of 3 or more years since patient's last colonoscopy, documented:
  - a. If Interval of 3 or more years since patient's last colonoscopy, documented equals Yes, include in Data Completeness Met and Performance Met.
    - Data Completeness Met and Performance Met letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a equals 30 procedures in the Sample Calculation.
  - b. If Interval of 3 or more years since patient's last colonoscopy, documented equals No, proceed

to check Documentation of medical reason(s) for an interval of less than 3 years since the last colonoscopy.

- 9. Check Documentation of medical reason(s) for an interval of less than 3 years since the last colonoscopy:
  - a. If Documentation of medical reason(s) for an interval of less than 3 years since the last colonoscopy equals Yes, include in Data Completeness Met and Denominator Exception.
    - Data Completeness Met and Denominator Exception letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter b<sup>1</sup> equals 10 procedures in the Sample Calculation.
  - b. If Documentation of medical reason(s) for an interval of less than 3 years since the last colonoscopy equals No, proceed to check Documentation of system reason(s) for an interval of less than 3 years since the last colonoscopy.
- 10. Check Documentation of system reason(s) for an interval of less than 3 years since the last colonoscopy:
  - a. If Documentation of system reason(s) for an interval of less than 3 years since the last colonoscopy equals Yes, include in the Data Completeness Met and Denominator Exception.
    - Data Completeness Met and Denominator Exception letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter b<sup>2</sup> equals 10 procedures in the Sample Calculation.
  - b. If Documentation of system reason(s) for an interval of less than 3 years since the last colonoscopy equals No, proceed to check Interval of less than 3 years since patient's last colonoscopy, reason not otherwise specified.
- 11. Check Interval of less than 3 years since patient's last colonoscopy, reason not otherwise specified:
  - a. If Interval of less than 3 years since patient's last colonoscopy, reason not otherwise specified equals Yes, include in Data Completeness Met and Performance Not Met.
    - Data Completeness Met and Performance Not Met letter is represented in the Data Completeness in the Sample Calculation listed at the end of this document. Letter c equals 20 procedures in the Sample Calculation.
  - b. If Interval of less than 3 years since patient's last colonoscopy, reason not otherwise specified equals

No, proceed to check Data Completeness Not Met.

- 12. Check Data Completeness Not Met:
  - If *Data Completeness Not Met*, the Quality Data Code or equivalent not submitted. 10 procedures have been subtracted from the Data Completeness Numerator in the Sample Calculation.

#### Sample Calculations

Data Completeness equals Performance Met (a equals 30 procedures) plus Denominator Exceptions (b<sup>1</sup> plus b<sup>2</sup> equals 20 procedures) plus Performance Not Met (c equals 20 procedures) divided by Eligible Population/Denominator (d equals 80 procedures). All equals 70 procedures divided by 80 procedures. All equals 87.5 percent.

Performance Rate equals Performance Met (a equals 30 procedures) divided by Data Completeness Numerator (70 procedures) minus Denominator Exceptions ( $b^1$  plus  $b^2$  plus 20 procedures). All equals 30 procedures divided by 50 procedures. All equals 60 percent.

\* See the posted measure specification for specific coding and instructions to submit this measure.

#### NOTE: Submission Frequency: Procedure

The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.